

CHAPTER IX

DES BÉNÉFICES DÉPENDANTS DE LA PROBABILITÉ DES ÉVÉNEMENTS FUTURS.

Pierre Simon Laplace

Théorie Analytique des Probabilités §§38–40, pp. 428–440

ON THE BENEFITS DEPENDING ON THE PROBABILITY OF FUTURE EVENTS.

If one awaits any number of simple events of which the probabilities are known and of which the arrival procures an advantage, their non-arrival causing a loss, to determine the mathematical benefit resulting from their awaiting. Expression of the probability that the real benefit will be comprehended within some given limits, when the number of events awaited is very great. However little advantage that each awaited event produces, the benefit becomes infinitely great and certain when the number of events is supposed infinite. N° 38.

If the diverse chances of an awaited event produce the advantages and the losses of which the respective probabilities are given, to determine the mathematical benefit resulting from the awaiting of any number of similar events. Expression of the probability that the real benefit will be comprehended within some given limits, when this number is very great. N° 39.

On the benefits of the establishments based on the probabilities of life. Expression of the capital that it is necessary to give in order to constitute a life pension on one or many heads. Expression of the rent that one individual must give to an establishment in order to assure to his heirs a capital payable at his death. Expression of the probability that the real benefit of the establishment will be comprehended within some given limits, by supposing that a great number of individuals, in constituting each a pension on his head, deposits each a determined sum into the funds of the establishment, in order to defray his expenses. N° 40.

38. We imagine that the arrival of an event procures the benefit ν , and that its non-arrival causes the loss μ . A person A awaits the arrival of a number s of similar events, all equally probable, but independent of one another; one demands what is his advantage.

Let q be the probability of the arrival of each event and consequently $1 - q$ that of its non-arrival; if one develops the binomial $[q + (1 - q)]^s$, the term

$$\frac{1.2.3\dots s}{1.2.3\dots i.1.2.3\dots (s-i)} q^i (1-q)^{s-i}$$

of this development will be the probability that out of s events i will arrive. In this case, the benefit of A is $i\nu$, and his loss is $(s-i)\mu$; the difference is $i(\nu + \mu) - s\mu$; by multiplying it by its probability expressed by the preceding term and taking the sum of these products for all the values of i , one will have the advantage of A, which, consequently, is equal to

$$-s\mu[q + (1-q)]^s + (\nu + \mu)S \frac{1.2.3\dots s}{1.2.3\dots i.1.2.3\dots (s-i)} q^i (1-q)^{s-i},$$

the sign S extending to all the values of i . One has

$$\begin{aligned} S \frac{i.1.2.3\dots s}{1.2.3\dots i.1.2.3\dots (s-i)} q^i (1-q)^{s-i} \\ = \frac{d}{dt} S \frac{1.2.3\dots s}{1.2.3\dots i.1.2.3\dots (s-i)} q^i t^i (1-q)^{s-i} = \frac{d}{dt} [qt + (1-q)]^s, \end{aligned}$$

provided that one supposed $t = 1$, after the differentiation, this which reduces this last member to qs ; the advantage of A is therefore $s[q\nu - (1-q)\mu]$. This advantage is null, if $\nu q = \mu(1-q)$, that is to say, if the benefit of the arrival of the event, multiplied by its probability, is equal to the loss caused by its non-arrival, multiplied by its probability. The advantage becomes negative and is changed into disadvantage, if the second product surpasses the first. In all cases, the advantage or the disadvantage of A is proportional to the number s of the events.

One will determine by the analysis of n° 16 the probability that the real benefit of A will be comprehended within the given limits, if s is a large number. Following this analysis, the sum of the diverse terms of the binomial $[q + (1-q)]^s$ comprehended between the two terms distant by $l + 1$, on both sides of the greatest, is

$$\frac{2}{\sqrt{\pi}} \int dt e^{-t^2} \frac{1}{\sqrt{2s\pi q(1-q)}} e^{-\frac{t^2}{2sq(1-q)}},$$

the integral being taken from $t = 0$ to $t = \frac{l}{\sqrt{2sq(1-q)}}$. The exponent of q in the greatest term is to very nearly, by the same section, equal to sq , and the exponents of q , corresponding to the extreme terms comprehended within the

preceding interval, are respectively $sq - l$ and $sq + l$. The benefits corresponding to these three terms are

$$\begin{aligned} & s[q\nu - (1 - q)\mu] - l(\nu + \mu), \\ & s[q\nu - (1 - q)\mu], \\ & s[q\nu - (1 - q)\mu] + l(\nu + \mu); \end{aligned}$$

by making therefore $l = r\sqrt{s}$, the probability that the real benefit of A will not be exceeded by the limits $s[q\nu - (1 - q)\mu] \pm r\sqrt{s}(\nu + \mu)$ is equal to

$$\frac{2}{\sqrt{\pi}} \frac{\int dr c^{-\frac{r^2}{2q(1-q)}}}{\sqrt{2q(1-q)}} + \frac{1}{\sqrt{2s\pi q(1-q)}} c^{-\frac{r^2}{2q(1-q)}},$$

the integral being taken from $r = 0$, and the last term being able to be neglected. One sees by this formula that if $q\nu - (1 - q)\mu$ is not null, the real benefit increases without ceasing and becomes infinitely great and certain in the case of an infinite number of events.

One is able to extend, by the following analysis, this result to the case where the probability of the s events are different, thus as the benefits and the losses which are attached. Let $q, q^{(1)}, q^{(2)}, \dots, q^{(s-1)}$ be the respective probabilities of these events; $\nu, \nu^{(1)}, \nu^{(2)}, \dots, \nu^{(s-1)}$ the benefits which their arrivals procure. One is able, in order to simplify, to set aside the losses which their non-arrivals cause, by comprehending in the benefit which the arrival of each event procures the quantity that A would lose by its non-arrival, and by subtracting next from the total advantage of A the sum of these last quantities; because it is easy to see that that changes not at all the position of A.

This posed, we will consider the product

$$\left(1 - q - qc^{\nu\varpi\sqrt{-1}}\right) \left(1 - q^{(1)} - q^{(1)}c^{\nu^{(1)}\varpi\sqrt{-1}}\right) \dots \left(1 - q^{(s-1)} - q^{(s-1)}c^{\nu^{(s-1)}\varpi\sqrt{-1}}\right).$$

It is clear that the probability that the sum of the benefits will be $f + l'$ will be equal to the coefficient of $c^{(f+l')\varpi\sqrt{-1}}$ in the development of this product; it is therefore equal to

$$(a) \quad \frac{1}{2\pi} \int d\varpi c^{-(f+l')\varpi\sqrt{-1}} \left(1 - q - qc^{\nu\varpi\sqrt{-1}}\right) \dots \left(1 - q^{(s-1)} - q^{(s-1)}c^{\nu^{(s-1)}\varpi\sqrt{-1}}\right).$$

the integral being taken from $\varpi = -\pi$ to $\varpi = \pi$, and the numbers $\nu, \nu^{(1)}, \dots$ being supposed, as one is able to make it, of whole numbers. We take the logarithm of the product

$$(b) \quad c^{-f\varpi\sqrt{-1}} \left(1 - q - qc^{\nu\varpi\sqrt{-1}}\right) \dots \left(1 - q^{(s-1)} - q^{(s-1)}c^{\nu^{(s-1)}\varpi\sqrt{-1}}\right);$$

by developing it according to the powers of ϖ , it becomes

$$(\text{S}q^{(i)}\nu^{(i)} - f)\varpi\sqrt{-1} - \frac{\varpi^2}{2}\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2} - \dots$$

the sign S reporting to all the values of i , from $i = 0$ to $i = s - 1$. The supposition of f equal to $\text{S}q^{(i)}\nu^{(i)}$ makes the first power of ϖ disappear, and the size of s , a very great number, renders insensible the terms depending on the powers of ϖ , superior to the square. By passing again therefore from the logarithms to the numbers in the preceding development, the product (b) becomes to very nearly

$$c^{-\frac{\varpi^2}{2}\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}},$$

this which changes the integral (a) into that here

$$\frac{1}{2\pi} \int d\varpi c^{-l'\varpi\sqrt{-1} - \frac{\varpi^2}{2}\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}}.$$

The integral must be taken from $\varpi = -\pi$ to $\varpi = \pi$, and $\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}$ being a great number of order s , it is clear that this integral is able to be extended without sensible error to the infinite positive and negative values of ϖ . By making therefore

$$\varpi \sqrt{\frac{\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}}{2}} + \frac{l'\sqrt{-1}}{\sqrt{2\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}}} = t$$

and integrating from $t = -\infty$ to $t = \infty$, the integral (a) becomes

$$\frac{1}{\sqrt{2\pi\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}}} c^{-\frac{l'^2}{2\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}}}.$$

If one multiplies this quantity by $2dl'$, and if next one integrates it from $l' = 0$, this integral will be the expression of the probability that the benefit of A will be comprehended within the limits $f \pm l'$, or $\text{S}q^{(i)}\nu^{(i)} \pm l'$; by making thus

$$l' = \sqrt{2\text{S}q^{(i)}(1 - q^{(i)})\nu^{(i)2}},$$

the probability that the benefit of A will be comprehended within the limits

$$Sq^{(i)}\nu^{(i)} \pm r\sqrt{2Sq^{(i)}(1-q^{(i)})\nu^{(i)2}}$$

is

$$\frac{2}{\sqrt{\pi}} \int dr e^{-r^2},$$

the integral being taken from $r = 0$.

Now it is necessary, by that which precedes, to change within the preceding limits $\nu^{(i)}$ into $\nu^{(i)} + \mu^{(i)}$ and to subtract it from $S\mu^{(i)}$; the probability that the real benefit of A will be comprehended within the limits

$$S[q^{(i)}\nu^{(i)} - (1-q^{(i)})\mu^{(i)}] \pm r\sqrt{2Sq^{(i)}(1-q^{(i)})(\nu^{(i)} + \mu^{(i)})^2}$$

is therefore

$$\frac{2}{\sqrt{\pi}} \int dr e^{-r^2}.$$

One sees by this formula that, as little as the mathematical expectation of each event surpasses zero, by multiplying the events to infinity, the first term of the expression of the limits being of order s , while the second is only of order \sqrt{s} , the real benefit is increased without ceasing and becomes at the same time infinitely great and certain, in the case of an infinite number of events.

39. We consider now the case where, at each event, the person A has any number of chances to hope or to fear. We suppose, for example, that an urn contains some balls of diverse colors, that one draws a ball from this urn, by replacing it into the urn after the drawing, and that the benefit of A is ν if the extracted ball is of the first color, that it is ν' if the extracted ball is of the second color, that it is ν'' if the extracted ball is of the third color, and thus consecutively, the benefits becoming negative when A is forced to give instead of receive. We name a, a', a'', \dots the probabilities that the ball extracted at each drawing will be of the first, or of the second, or of the third, etc. color, and we suppose that one has thus s drawings; one will have first

$$a + a' + a'' + \dots = 1.$$

By multiplying next the terms of the first member of this equation, respectively by $c^{\nu\varpi\sqrt{-1}}$, $c^{\nu'\varpi\sqrt{-1}}$, $c^{\nu''\varpi\sqrt{-1}}$, ..., the term independent of the powers of $c^{\varpi\sqrt{-1}}$, in the development of the function

$$c^{-(l+s\mu)\varpi\sqrt{-1}} \left(ac^{\nu\varpi\sqrt{-1}} + a'c^{\nu'\varpi\sqrt{-1}} + a''c^{\nu''\varpi\sqrt{-1}} + \dots \right)^s,$$

will be, by that which precedes, the probability that, in s drawings, the benefit of A will be $s\mu + l$; this probability is therefore equal to

$$(c) \quad \frac{1}{2\pi} \int d\varpi c^{-l\varpi\sqrt{-1}} \left[c^{-\mu\varpi\sqrt{-1}} \left(c^{\nu\varpi\sqrt{-1}} + a'c^{\nu'\varpi\sqrt{-1}} + \dots \right) \right]^s,$$

the integral relative to ϖ being taken from $\varpi = -\pi$ to $\varpi = \pi$. If one develops with respect to the powers of ϖ the hyperbolic logarithm of the quantity raised to the power s under the \int sign, and if one observes that $a + a' + a'' + \dots = 1$, one will have, for this logarithm,

$$\begin{aligned} & \varpi\sqrt{-1}(a\nu + a'\nu' + a''\nu'' + \dots - \mu) \\ & - \frac{\varpi^2}{2} [a\nu^2 + a'\nu'^2 + a''\nu''^2 + \dots - (a\nu + a'\nu' + a''\nu'' + \dots)^2 - \dots \end{aligned}$$

One will make the first power of ϖ disappear, by making

$$\mu = a\nu + a'\nu' + a''\nu'' + \dots;$$

if one supposes next

$$2k^2 = a\nu^2 + a'\nu'^2 + a''\nu''^2 + \dots - (a\nu + a'\nu' + a''\nu'' + \dots)^2,$$

and if one observes that, s being supposed a great number, one is able to neglect the powers of ϖ superior to the square, one will have, by passing again from the logarithms to the numbers,

$$\left[c^{-\mu\varpi\sqrt{-1}} \left(ac^{\nu\varpi\sqrt{-1}} + a'c^{\nu'\varpi\sqrt{-1}} + a''c^{\nu''\varpi\sqrt{-1}} + \dots \right) \right]^s = c^{-sk^2\varpi^2},$$

this which changes the integral (c) into that here

$$\frac{1}{2\pi} \int d\varpi c^{-l\varpi\sqrt{-1} - sk^2\varpi^2},$$

which becomes, by integrating as in the preceding section,

$$\frac{1}{2k\sqrt{s\pi}} e^{-\frac{l^2}{4sk^2}}.$$

By multiplying it by $2dl$ and integrating the product from $l = 0$, one will have the probability that the real benefit of A will be comprehended within the limits

$$s(a\nu + a'\nu' + a''\nu'' + \dots) \pm l;$$

by making therefore

$$l = 2kr'\sqrt{s},$$

this probability will be, by taking the integral from $r' = 0$,

$$\frac{2}{\sqrt{\pi}} \int dr' e^{-r'^2}.$$

We have supposed in that which precedes the probabilities of the events known; we examine the case where they are unknown. We suppose that, out of m similar events awaited, n are arrived, and that A awaits s similar events, of which each procures to him by its arrival the benefit ν , the non-arrival causing to him the loss μ . If one represents by $\frac{n}{m}s + z$ the number of events which will arrive out of the s events awaited, the probability that z will be contained within the limits $\pm kt$ will by n° 30

$$\frac{2}{\sqrt{\pi}} \int dt e^{-t^2},$$

the integral being taken from $t = 0$, k^2 being equal to

$$\frac{2ns(m-n)(m+s)}{m^3}.$$

But, $\frac{n}{m}s + z$ being the number of arrived events, the real benefit of A is

$$\left[\frac{n\nu}{m} - \frac{(m-n)\mu}{m} \right] s + z(\nu + \mu);$$

the preceding integral is therefore the probability that the real benefit of A will be comprehended within the limits

$$\left[\frac{n\nu}{m} - \frac{(m-n)\mu}{m} \right] s + kt(\nu + \mu).$$

k is of order \sqrt{s} , if m and n are of order equal or greater than s ; thus, howsoever small that the mathematical expectation be relative to each event, the real benefit becomes at infinity, certain and infinitely great, when the number of past events is supposed infinite, as the one of future events.

40. We will now determine the benefits of the establishments founded on the probabilities of human life. The most simple way to calculate these benefits is to reduce them to actual capitals. We take for example life pensions. A person of age A wishes to constitute on his head a life pension h ; one demands the capital that he must give for it to the funds of the establishment which makes this pension for him.

If one names y_0 the number of individuals of age A in the Table of mortality of which one makes use, and y_x the number of individuals of age $A + x$, the probability to pay the pension at the end of the year $A + x$ will be $\frac{y_x}{y_0}$; consequently, the value of the payment will be $\frac{hy_x}{y_0}$. But, if one designates by r the annual interest on unity, so that the capital 1 becomes $1 + r$ after one year, it will become $(1 + r)^x$ after x years; thus, the payment $(1 + r)^x$ made at the end of the x^{th} year, reduced to actual capital, becomes unity, or this same payment divided by $(1 + r)^x$; the payment $\frac{hy_x}{y_0}$ reduced to actual capital is therefore $\frac{hy_x}{y_0(1+r)^x}$. The sum of all the payments made during the duration of life of the person who constitutes the pension and multiplied by their probability is equivalent therefore to an actual capital represented by the finite integral

$$\sum \frac{hy_x}{y_0(1+r)^x},$$

the characteristic \sum must embrace all the values of the function that it affects.

One is able to determine this integral by forming all these values according to the Table of mortality, and by adding them together; one will deduce next the capitals from one another, by observing that, if one names F the capital relative to the age A and F' the capital relative to age $A + 1$, one has

$$F = \frac{y_1}{y_0} \frac{F' + h}{1 + r}.$$

But this process is simplified when the law of mortality is known, and especially when it is given by a rational and entire function of x , this which is always possible, by considering the numbers of the Table of mortality as some ordinates of which the corresponding ages are the abscissas, and by making pass a parabolic curve through the extremities of the two extreme ordinates and of many intermediate ordinates. The differences which exist among the diverse Tables of mortality permit regarding this mean as exact also as these Tables, and even to be satisfied with a small number of ordinates.

We make

$$\frac{1}{1+r} = p, \quad \frac{y_x}{y_0} = u;$$

we resume formula (16) of n° 11 of Book I which gives

$$\sum p^x u = \frac{p^x}{p^{\frac{du}{dx}} - 1} + f,$$

f being an arbitrary constant. It is necessary, in the development of the first term of the second member of this equation with respect to the ratio to the powers of $\frac{du}{dx}$, to change any power $\left(\frac{du}{dx}\right)^i$ into $\frac{d^i u}{dx^i}$, and to multiply by u the first term, which is independent of $\frac{du}{dx}$. One has thus

$$\sum p^x u = f - \frac{p^x u}{1-p} - \frac{p^{x+1} \frac{du}{dx}}{(1-p)^2} - \frac{(p+1)p^{x+1}}{1.2.(1-p)^3} \frac{d^2 u}{dx^2} + \dots$$

In order to determine f , one will observe that the integral $\sum p^x u$ is null when $x = 1$, and that it is terminated when $x = n + 1$, $A + n$ being the limit of life; because then it embraces the terms corresponding to all the numbers 1, 2, 3, ..., n . We designate therefore by (u) , $\left(\frac{du}{dx}\right)$, ..., u' , $\left(\frac{du'}{dx}\right)$, ... the values of u , $\frac{du}{dx}$, ..., corresponding to $x = 1$ and to $x = n + 1$; one will have

$$(o) \quad \sum \frac{h p^x y_x}{y_0} = h \left\{ \begin{array}{l} \frac{p}{1-p} [(u) - p^n (u')] \\ + \frac{p^2}{(1-p)^2} \left[\left(\frac{du}{dx}\right) - p^n \left(\frac{du'}{dx}\right) \right] \\ + \frac{(p+1)p^2}{1.2.(1-p)^3} \left[\left(\frac{d^2 u}{dx^2}\right) - p^n \left(\frac{d^2 u'}{dx^2}\right) \right] \\ + \dots \end{array} \right\}.$$

If u or $\frac{y_x}{y_0}$ is constant and equal to unity, from $x = 1$ to $x = n$, then the life pension must be paid certainly during the number n of years, and it becomes an annuity. In this case, $\frac{du}{dx}$ is null, and the preceding formula gives $\frac{hp(1-p^n)}{1-p}$ for the capital equivalent to the annuity h .

If $u = 1 - \frac{x}{n}$, then the probability of life decreases in arithmetic progression, and the preceding formula gives

$$\frac{hp}{1-p} \left[1 - \frac{1-p^n}{n(1-p)} \right]$$

for the capital equivalent to the life annuity h , and thus consecutively.

We suppose now that one wishes to constitute a life pension h on many individuals of ages $A, A + a, A + a + a', \dots$, so that the pension remains to the survivors. We designate by $y_x, y_{x+a}, y_{x+a+a'}, \dots$ the numbers of the Table of mortality, corresponding to the ages $A, A + a, A + a + a', \dots$, the probability that the first individual has to live to the age $A + x$ being $\frac{y_x}{y_0}$, the probability that at this age he will have ceased to live is $1 - \frac{y_x}{y_0}$. Similarly, the probability that the second individual has to live to age $A + a + x$ or to the end of the x^{th} year of the constitution of the pension being $\frac{y_{x+a}}{y_a}$, the probability that he will have ceased to live then is $1 - \frac{y_{x+a}}{y_a}$; the probability that the third individual will have ceased to live, at the same epoch of the constitution of the pension, is $1 - \frac{y_{x+a+a'}}{y_{a+a'}}$, and thus consecutively. The probability that none of these individuals will exist at this epoch is

$$\left(1 - \frac{y_x}{y_0} \right) \left(1 - \frac{y_{x+a}}{y_a} \right) \left(1 - \frac{y_{x+a+a'}}{y_{a+a'}} \right) \dots$$

By subtracting this product from unity, the difference will be the probability that one of these individuals at least will be living at the end of the x^{th} year of the constitution of the pension. We name u this probability; $\sum hp^x u$ will be the actual capital equivalent to the life annuity h . But one must observe, by taking this integral, that the quantities y_x, y_{x+a}, \dots are nulls, when their indices $x, x + a, \dots$ surpasses the number $n, A + n$ being the limit of the life.

If y_x is a rational and entire function of x , and of exponentials such as q^x, r^x, \dots one will have easily, by the formulas of Book I, the integral $h \sum p^x u$; but one is able in all cases to form, by means of a Table of mortality, all the terms of this integral, by taking the sum and constructing thus some Tables of life annuities on one or many heads.

The preceding analysis similarly serves to determine the life annuity that one must make at an establishment in order to assure to his heirs a capital after his death. The capital equivalent to the life annuity h , made on a person of age A , is, by that which precedes, $hS \frac{p^x y_x}{y_0}$, the sign S comprehending all the terms inclusively, from $x = 1$ to the limit of the life of the person. We name hq this integral, and we imagine that the establishment receives from this person the rent h , and gives to him in exchange the capital hq . We imagine next that the same person places this capital at perpetual interest with the establishment itself, the annual interest of unity being r or $\frac{1-p}{p}$. It is clear that the establishment must render the capital hq to the heirs of the person. But the person has made during his life the rent h to the establishment, and the person has received from it the pension $\frac{hq(1-p)}{p}$; the rent that the person has made really is therefore $h \left[1 - \frac{q(1-p)}{p} \right]$; it is therefore this that the person must give annually to the establishment in order to assure to his heirs the capital hq .

I will not insist further on these objects, in the same way with respect to those which are relative to the establishments of assurance of each kind, because they present no difficulty. I will observe only that all these establishments must, in order to prosper, reserve themselves a benefit and multiply considerably their affairs, so that, their real benefit becoming near certain, they are exposed the least that it is possible to some great losses which would be able to destroy them. In effect, if the number of the affairs is s and if the advantage of the establishment in each of them is b , then it becomes extremely probable that the real benefit of the establishment will be sb , s being supposed a very great number.

In order to show it, we will suppose that s persons of age A constitute, each on his head, a life annuity h , and we will consider one of these persons who we will designate by C . If C dies in the interval at the end of the year x elapsed from the constitution of his pension to the end of the year $x + 1$, the establishment will have paid to him the pension h during x years, and the sum of these payments, reduced to actual capital, will be $h(p+p^2 + \dots + p^x)$ or $\sum hp^{x+1}$; now the probability that C will die within this interval is $\frac{y_x - y_{x+1}}{y_0}$ or $-\frac{\Delta y_x}{y_0}$; the value of the loss that the establishment must then support is therefore $-\frac{\Delta y_x}{y_0} \sum hp^{x+1}$. The sum of all these losses is

$$(r) \quad - \sum \left(\frac{\Delta y_x}{y_0} \sum hp^{x+1} \right);$$

this is the capital that C must deposit in the funds of the establishment in order to receive from it the life pension h . One is able to observe here that one has

$$-\Delta y_x \sum p^{x+1} = -y_{x+1} \sum p^{x+1} + y_x \sum p^x + y_x p^x;$$

by integrating the second member of this equation, the function (r) is reduced to

$$-\frac{\Delta y_x}{y_0} \sum h p^x + \frac{\sum h y_x p^x}{y_0} \text{const.};$$

now $\sum p^x$ is reduced to zero, when $x = 1$, and when $x = n + 1$ y_x is null by that which precedes; the function (r) or the capital that C must pay to the establishment is therefore $\frac{\sum h y_x p^x}{y_0}$, this which is conformed to that which precedes. But under the form of the function (r) one is able to apply to the benefit of the establishment the analysis of n° 39. In effect, one has in this case, by the section cited,

$$a\nu + a'\nu' + a''\nu'' + \dots = -\sum \left(\frac{\Delta y_x}{y_0} \sum h p^{x+1} \right);$$

next a, a', \dots being the successive values of $-\frac{\Delta y_x}{y_0}$, one will have

$$a\nu + a'\nu'^2 + \dots = \sum \left[-\frac{\Delta y_x}{y_0} (\sum h p^{x+1})^2 \right],$$

so that

$$2k^2 = \sum \left[-\frac{\Delta y_x}{y_0} (\sum h p^{x+1})^2 \right] - \left[\sum \frac{\Delta y_x}{y_0} (\sum h p^{x+1}) \right]^2.$$

In supposing that each of s persons who constitute the pension h on his head deposit to the funds of the establishment, beyond the capital corresponding to this pension, a sum b , in order to defray the expense of the establishment, one will have, by n° 39,

$$\frac{2}{\sqrt{\pi}} \int dr' c^{-r'^2},$$

for the probability that the real benefit of the establishment will be comprehended within the limits

$$sb \pm 2kr' \sqrt{s}.$$

Thus, in the case of an infinite number of affairs, the real benefit of the establishment becomes certain and infinite. But then those who treat with it have a mathematical disadvantage which must be compensated by a moral advantage, of which the estimation will be the object of the following Chapter.

© 2007 Richard J. Pulskamp
All rights reserved.